

DIBOSON PRODUCTION CROSS SECTIONS AT $\sqrt{s} = 1.96$ TeV

J. SEKARIC

*Department of Physics, 513 Keen Building,
Florida State University, Tallahassee, Florida, USA*

for the CDF and DØ Collaborations

The increasing size of the data recorded by the CDF and DØ experiments at the Tevatron collider at $\sqrt{s} = 1.96$ TeV makes the diboson physics program more accessible for probes of the electroweak gauge structure in the Standard Model. Here we summarize the most recent measurements of the diboson cross sections and limits on the trilinear gauge boson couplings.

1 Introduction

The diboson production plays an important role in electroweak precision measurements and searches for the New Physics (NP) which may exist at some energy scale Λ . The search for the NP is often related to the precision measurements such those of the cross sections and the trilinear gauge boson couplings (TGCs). More discussions about these TGC parameters can be found in¹. Depending on the NP scenario, these observables are expected to deviate from their SM predictions. The charged TGCs studied in WW and $W\gamma$ production are $(\Delta g_1^Z, \Delta \kappa_{\gamma, Z}, \lambda)$ and $(\Delta \kappa_\gamma, \lambda)$, respectively, and $(\Delta g_1^Z, \Delta \kappa_Z, \lambda)$ in WZ production, where Δ represents the deviation from the SM prediction. In the SM, $\Delta g_1^Z = \Delta \kappa_{\gamma, Z} = \lambda = 0$. The neutral TGCs $h_{30,40}^{\gamma, Z}$ studied in $Z\gamma$ production are not allowed in the SM and their values are predicted to be zero. Besides, the diboson production is an important background to studies of the top quark and searches for the Higgs boson and SUSY particles. The most precise knowledge of background processes and their proper modeling is highly valuable for current and future studies.

2 Diboson Production

2.1 $Z\gamma \rightarrow \nu\nu\gamma$

The $Z\gamma$ events are reconstructed from 3.6 fb^{-1} of DØ data. Candidate events are required to have one isolated central photon ($|\eta_{det}| < 1.1$ ²) with $E_T > 90$ GeV and $\cancel{E}_T > 70$ GeV. The pointing algorithm³ is used in order to reduce the contribution from bremsstrahlung photons. After all selection criteria were applied 51 $\nu\nu\gamma$ candidate events are observed. The predicted numbers of signal and background events are 33.7 ± 3.4 and 17.3 ± 2.4 , respectively. The dominant background events are $W \rightarrow e\nu$ in which the electron is misidentified as a photon and it contributes with 9.7 ± 0.6 events. The measured cross section is $\sigma_{ZZ} \times BR(Z \rightarrow \nu\nu) = 32 \pm 9(\text{stat} + \text{syst}) \pm 2(\text{lumi}) \text{ fb}$ ⁵ which is in agreement with the next-to-leading (NLO) cross section of $(39 \pm 4) \text{ fb}$ ⁴. The observed signal significance is 5.1 standard deviations (s.d.).

Furthermore, the photon E_T spectrum shown in Fig. 1, is used to set the limits on $ZZ\gamma$ and $Z\gamma\gamma$ TGCs. The 95% C.L one-dimensional limits for $h_{30,40}^{\gamma,Z}$ at $\Lambda = 1.5$ TeV are $|h_{30}^{\gamma}| < 0.036$, $|h_{30}^Z| < 0.035$ and $|h_{40}^{\gamma,Z}| < 0.0019$ ⁵. The combination with the previous results in the most restrictive limits on these couplings at 95% C.L. of $|h_{30}^{\gamma,Z}| < 0.033$ and $|h_{40}^{\gamma,Z}| < 0.0017$ of which three of them (h_{40}^{γ} , h_{40}^Z and h_{30}^Z) are world's best to date.

2.2 $ZZ \rightarrow \nu\nu l^+ l^-$

For the first time, the ZZ production in the $\nu\nu ll$ ($l = e, \mu$) final states has been studied at the DØ using 2.7 fb^{-1} of data⁶. The analysis builds up a new variable $\cancel{E}'_T$, highly discriminating against the $Z \rightarrow ll$ background events. Its purpose is to minimize the mismeasurement of the transverse momentum of either the charged leptons or the hadronic recoil system which contributes to the reconstructed $\cancel{E}_T$. In the electron channel $\cancel{E}'_T$ is required to be > 27 GeV, and in the muon channel $\cancel{E}'_T > 30(35)$ GeV for data collected during 2002-2006 (2006-2008). In addition, each channel requires that there are only two oppositely charged isolated leptons with $p_T > 15$ GeV with the dilepton invariant mass is $(70 < M_{ll} < 110)$ GeV. Events with additional lepton of the same family are vetoed. Further separation between the signal and WW , WZ , W +jets, $ZZ \rightarrow l^+ l^-$ backgrounds uses the likelihood discriminant. The total number of 43 candidate events has been selected of which 26.5 ± 0.5 are predicted to be the background. The observed signal significance is 2.6 s.d. and the measured cross section is $\sigma_{ZZ} = 2.01 \pm 0.93(stat) \pm 0.29(syst)$ pb which is consistent with the NLO SM predicted cross section of 1.4 ± 0.1 pb⁷.

2.3 $ZZ \rightarrow l^+ l^- l'^+ l'^-$

Sensitivity to single lepton cuts plays an important role when selecting $l^+ l^- l'^+ l'^-$ events ($l, l' = e \text{ or } \mu$). Application of tighter selection criteria such as lepton p_T , dilepton invariant mass and lepton isolation, relative to the previous DØ analysis⁸, results in the first observation of the ZZ production at a hadron collider. The analysis uses 1.7 fb^{-1} of DØ data in which four isolated leptons with $p_T > 15$ GeV are required to be within $|\eta_{det}| < 2$ if muons and $|\eta_{det}| < 1.1$ or $1.5 < |\eta_{det}| < 3.2$ if electrons. Two most energetic leptons in the $\mu\mu\mu\mu$ or $eeee$ channel are required to have $p_T > 30(25)$ GeV. Events are required to have at least one Z with an invariant mass greater than 70 GeV and the other greater than 50 GeV. In the $ee\mu\mu$ channel the two most energetic electrons or muons must have $p_T > 25(15)$ GeV. Lepton candidates are required to be spatially separated by $R > 0.2$ ($R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$) to reduce $Z \rightarrow \mu\mu$ background. The total of 3 events, two in the $eeee$ channel and one in the $\mu\mu\mu\mu$ channel, were observed. The SM signal and background are expected to contribute with 1.89 ± 0.08 and $0.14^{+0.03}_{-0.02}$ events, respectively. The observed signal significance is 5.3σ and the measured cross section $\sigma_{ZZ} = 1.75^{+1.27}_{-0.86}(stat) \pm 0.13(syst)$ pb⁹ is in agreement with the SM prediction⁷. The combination with previous analyses⁶ and ⁸ results in the observed signal significance of 5.7σ and the measured cross section is $\sigma_{ZZ} = 1.60 \pm 0.63(stat)^{+0.16}_{-0.17}(syst)$ pb.

2.4 $WZ \rightarrow jjll$

A search for anomalous ZWW TGCs in the $jjll$ final states results in setting the 95% C.L. limits on $\Delta\kappa_Z, \lambda$ and Δg_1^Z using the p_T^ll distribution, and the 95% C.L. limits on the cross section for the WZ production¹⁰. This preliminary result is obtained using 1.9 fb^{-1} of CDF data selecting events with two high p_T leptons and two jets. The p_T^ll phase space is divided in three regions: control region used to validate the data modeling (105 – 140 GeV), medium (140 – 210 GeV) and high (> 210 GeV) regions. The medium and high p_T^ll regions are used to perform the measurements. Total number of observed (predicted) events is 97 (71.4 ± 0.5) and

12 (9.74 ± 0.18) in the medium and high region, respectively. The observed 95% C.L. limits on TGCs are $-1.09 < \Delta\kappa_Z < 1.40$, $-0.18 < \lambda < 0.18$ and $-0.22 < \Delta g_1^Z < 0.32$ for $\Lambda = 1.5$ TeV, and $-1.01 < \Delta\kappa_Z < 1.27$, $-0.16 < \lambda < 0.17$ and $-0.20 < \Delta g_1^Z < 0.29$ for $\Lambda = 2$ TeV. Unbinned MC fit to data in the dijet mass distribution results in the 95% C.L. cross section limits of 234 fb and 135 fb for medium and high region, respectively.

2.5 $WW \rightarrow l\nu l'\nu$

The most precise WW cross section measurement at a hadron collider is performed analyzing the $l\nu l'\nu$ ($l, l' = e, \mu$) final states with 1.0 fb^{-1} of $D\bar{O}$ data¹¹. In each ll' final state ($ee, \mu\mu$ or $e\mu$) the two most energetic leptons are required to have $p_T > 25$ (15) GeV, to be of opposite charge and to be spatially separated from each other by $R > 0.8$ (ee) and $R > 0.5$ ($e\mu$). The $Z/\gamma^* \rightarrow ll$ background is effectively removed requiring $\cancel{E}_T > 45$ (ee), 20 ($e\mu$) or 35 ($\mu\mu$) GeV, $\cancel{E}_T > 50$ GeV if $|M_Z - m_{ee}| < 6$ GeV (ee), $\Delta\phi_{\mu\mu} < 2.45$ and $\cancel{E}_T > 40$ GeV if $\Delta\phi_{e\mu} > 2.8$. Imposing the upper cut on the transverse momentum of the WW system, of 20 (ee), 25 ($e\mu$) and 16 ($\mu\mu$) GeV minimizes the $t\bar{t}$ background. After all selection criteria were applied, all three combined channels yield 100 candidate events, 38.19 ± 4.01 predicted background events and 64.70 ± 1.12 predicted signal events. The cross section measurements in the individual channels are combined, yielding $\sigma_{WW} = 11.5 \pm 2.1(\text{stat} + \text{syst}) \pm 0.7(\text{lumi})$ pb which is in agreement with the SM NLO prediction of $12.4 \pm 0.8 \text{ pb}$ ⁷. The p_T distributions of the leading and trailing leptons were used to set limits on anomalous TGCs considering two different parameterizations between the couplings. The one-dimensional 95% C.L. limits for $\Lambda = 2$ TeV are $-0.54 < \Delta\kappa_\gamma < 0.83$, $-0.14 < \lambda_\gamma = \lambda_Z < 0.18$ and $-0.14 < \Delta g_1^Z < 0.30$ under the $SU(2)_L \times U(1)_Y$ -conserving constraints¹², and $-0.12 < \Delta\kappa_\gamma = \Delta\kappa_Z < 0.35$ and $-0.14 < \lambda_\gamma = \lambda_Z < 0.18$ under the assumption that γWW and ZWW couplings are equal.

2.6 $WW + WZ \rightarrow l\nu jj$

This analysis results in the first evidence for the WW/WZ production in the $l\nu jj$ final states at the hadron collider with 1.1 fb^{-1} of $D\bar{O}$ data¹³. Selected $l\nu jj$ ($l = e, \mu$) candidate events are required to have a single isolated lepton with $p_T > 20$ GeV and $|\eta| < 1.1$ (2.0) for electrons (muons), $\cancel{E}_T > 20$ GeV and at least two jets with $p_T > 20$ GeV. The jet of highest p_T must have $p_T > 30$ GeV and the transverse mass of leptonically decaying W boson must be > 35 GeV to reduce the multijet background. Because of the small signal-to-background ratio (3%), an accurate modeling of the dominant W +jets background is essential and therefore, studied in great detail. After all selection criteria were applied, the signal and the backgrounds are further separated using a multivariate classifier, Random Forest (RF)¹⁴. The signal cross section is determined from a fit of signal and background RF templates to the data with respect to variations in the systematic uncertainties¹⁵ and is measured to be $\sigma_{WW+WZ} = 20.2 \pm 2.5(\text{stat}) \pm 3.6(\text{syst}) \pm 1.2(\text{lumi})$ pb which is consistent with the SM NLO prediction of $\sigma(WW + WZ) = 16.1 \pm 0.9 \text{ pb}$ ⁷. The observed signal significance is 4.4 s.d.. Fig. 2 shows the WW/WZ dijet mass peak extracted from data compared to the MC prediction.

3 Summary

The recent results in diboson production at the Tevatron, using $1.0 - 3.6 \text{ pb}^{-1}$ of data, have been presented. Measured cross sections for the $Z\gamma, ZZ, WZ$ and WW processes and TGCs at the $\gamma ZZ, \gamma WW$ and ZWW vertices are in agreement with the SM predictions. The $D\bar{O}$ experiment sets the world's tightest limits on the h_{40}^γ, h_{40}^Z and h_{30}^Z TGCs and reports the first evidence of WW/WZ production in semi-leptonic final states at a hadron collider.




Figure 1: Photon E_T spectrum of $\nu\nu\gamma$ data candidate events compared to the SM signal and background, and the expected distribution in the presence of anomalous TGCs. The systematic and statistical uncertainties on the SM MC events are included as shaded bands.




Figure 2: A comparison of the extracted signal (filled histogram) to background-subtracted data (points), along with the ± 1 standard deviation (s.d.) systematic uncertainty on the background. The residual distance between the data points and the extracted signal, divided by the total uncertainty, is given at the bottom.

References

1. K. Hagiwara, J. Woodside, and D. Zeppenfeld, *Phys. Rev. D* **41**, 2113 (1990).
2. DØ uses a cylindrical coordinate system with the z axis running along the beam axis. Angles θ and ϕ are the polar and azimuthal angles, respectively. Pseudorapidity is defined as $\eta = -\ln[\tan(\theta/2)]$ in which θ is measured with respect to the primary vertex. In the massless limit, η is equivalent to the rapidity $y = (1/2) \ln[(E + p_z)/(E - p_z)]$. η_{det} is the pseudorapidity measured with respect to the center of the detector.
3. S. Kesisoglou, *FERMILAB-THESES-2004*, UMI-31-74625 (2004).
4. U. Baur, T. Han and J. Ohnemus, *Phys. Rev. D* **57**, 2823 (1998).
5. The DØ Collaboration: V. M. Abazov *et al*, FERMILAB-PUB-09-046-E, Accepted by *Phys. Rev. Lett.* (2009).
6. The DØ Collaboration: V. M. Abazov *et al*, *Phys. Rev. D* **78**, 072002 (2008).
7. J. M. Campbell and R.K. Ellis, *Phys. Rev. D* **60**, 113006 (1999). Cross sections were calculated with the same parameter values given in the paper, except with $\sqrt{s} = 1.96$ TeV.
8. The DØ Collaboration: V. M. Abazov *et al*, *Phys. Rev. Lett.* **101**, 131801 (2008).
9. The DØ Collaboration: V. M. Abazov *et al*, *Phys. Rev. Lett.* **101**, 171803 (2008).
10. The CDF Collaboration: A. Abulencia *et al*, <http://www-cdf.fnal.gov/physics/ewk/2008/WWZaTGC/>
11. The DØ Collaboration: V. M. Abazov *et al*, FERMILAB-PUB-09-094-E, submitted to *Phys. Rev. Lett.* (2009).
12. M. Bilenky, J. L. Kneur, F. M. Renard and D. Schildknecht, *Nucl. Phys. B* **409**, 22 (1993).
13. The DØ Collaboration: V. M. Abazov *et al*, *Phys. Rev. Lett.* **102**, 161801 (2009).
14. L. Breiman, *Machine Learning* **45**, 5 (2001); I. Narsky, [arXiv:physics/0507143] [physics.data-an] (2005).
15. W. Fisher, FERMILAB-TM-2386-E (2006).